

Two-photon scattering of a tightly focused weak light beam from a small atomic ensemble

Dibyendu Roy

*Theoretical Division and Center for Nonlinear Studies,
Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

We study two-photon scattering of a tightly focused weak light beam from a small atomic ensemble of two-level atoms (2LAs). The scaling of the two-photon nonlinearity at single photon resonance shows a non-monotonic behaviour with increasing number of few identical 2LAs. Single-photon transport in a system of two 2LAs is similar to that in a V -type three-level atom (3LA). However two-photon transport in these two systems shows very different line-shapes. When the single-photon transmission is zero in these systems, two transmitted photons are bunched together in a V -type 3LA, while their correlation is zero in two 2LAs. The difference in the two-photon line-shape persists for few 2LAs and 3LAs. Therefore, the two-photon scattering of a tightly focused weak light beam can be used as a probe to detect atomic level structures of different atoms with similar transition energies.

A strong photon-photon interaction at the level of weak light field is one main challenge for realizing photonic quantum information devices. It is efficiently achieved by placing a nonlinear medium, such as an atom or a superconducting qubit in a cavity. The cavity greatly enhances the coupling between the medium and the photons. Recently a new approach to realize a strong two-photon nonlinearity at the level of few photons has been demonstrated using scattering of a tightly focused light beam from the dipole moment of an atom [1–4]. It can be realized easily by confining the light beam and the atom in a one-dimensional (1D) photonic waveguide [5–13]. Tight confinement of light fields in the waveguide directs the majority of the spontaneously emitted light from the atom into the transport channels, while local interactions at the atom induce a strong photon-photon correlation by preventing multiple occupancy of photons at the atom. A single two-level atom (2LA) is highly saturated by a single photon, thus it creates a strong optical nonlinearity for multiple incident photons. However an atomic ensemble can not be saturated by a single photon when the number of atoms is large, and the photon-photon interaction declines with an increasing number of 2LAs. We calculate a scaling of the two-photon nonlinearity with an increasing number of few identical 2LAs. The decay of two-photon nonlinearity shows a nonmonotonic behaviour with the increasing atom number at single photon resonance of a 2LA. However the two-photon nonlinearity is diminished monotonically with increasing atoms for incident photons detuned from single-photon resonance. We also derive two-photon scattering by two different side-coupled 2LAs [14, 15]. The single-photon scattering by two different side-coupled 2LAs in a 1D waveguide is similar to that by a V -type three level atom (3LA) [16]. However, the nature of two-photon scattering in these two systems is quite different. When the single-photon transmission is zero in these systems, two reflected photons are anti-bunched in a V -type 3LA, but they are not anti-bunched in two 2LAs. Two transmitted photons at these parameters are bunched in a 3LA while




FIG. 1. A schematic of a small ensemble of atoms in a tightly focused weak light beam.

there is no transmitted photon for the 2LAs. The difference in the two-photon line-shape persists for few 2LAs and V -type 3LAs. Therefore, we propose that the two-photon scattering line-shape of a tightly focused weak light beam can be used to probe atomic level structures of different type of atomic ensembles.

We consider N 2LAs side-coupled to the propagating left- and right-moving photon modes. These photon modes can be confined in a 1D waveguide [7] or they can be created tightly focused by any other experimental techniques [3]. For simplicity of calculation, we assume all the atoms are connected to the chiral photon modes at a same position. This is a valid approximation for a small ensemble of atoms, $R \ll \lambda$ where R is radius of the ensemble and λ is resonant radiation wavelength. The Hamiltonian of the full system,

$$\mathcal{H} = -iv_g \int dx [a_R^\dagger(x) \partial_x a_R(x) - a_L^\dagger(x) \partial_x a_L(x)] + \sum_{i=1}^N (\tilde{\Omega}_i a_{ei}^\dagger a_{ei} + [V(a_R^\dagger(0) + a_L^\dagger(0)) \sigma_{i-} + H.c.]) , \quad (1)$$

where $\tilde{\Omega}_i = \Omega_i - i\gamma_i/2$. Here v_g is the group velocity of photons, and $a_R^\dagger(x)$ ($a_L^\dagger(x)$) is a bosonic creation operator for a right (left)-moving photon at position x . We have set ground state energy of 2LAs to be zero, and Ω_i is




FIG. 2. (a) Two different two-level atoms and (b) a V-type three-level atom in a 1D photonic waveguide with incident, transmitted and reflected light beams.

energy of the excited state of i th 2LA. Here $a_{ei}^\dagger$ ($a_{gi}^\dagger$) is a creation operator of the i th 2LA's excited (ground) state, and $\sigma_{i-} = a_{gi}^\dagger a_{ei}$. The coupling strength between the photon modes and an 2LA is V . It is important to incorporate spontaneous emission from the atoms to the other modes apart from the left and right moving photon modes of the Hamiltonian in Eq.1. This is done here by including an imaginary term $-i\gamma_i/2$ in the energy of the excited atomic state within the quantum jump picture [17]. Next we transform the Hamiltonian using, $a_e^\dagger(x) \equiv \frac{1}{\sqrt{2}}(a_R^\dagger(x) + a_L^\dagger(-x))$ and $a_o^\dagger(x) \equiv \frac{1}{\sqrt{2}}(a_R^\dagger(x) - a_L^\dagger(-x))$ which decouple $\mathcal{H}$ in two parts, $\mathcal{H} = \mathcal{H}_e + \mathcal{H}_o$. One part $\mathcal{H}_e$ contains interaction of atoms with even photon modes while the other part $\mathcal{H}_o$ has only free propagating photons in odd modes. $\mathcal{H}_e = -iv_g \int dx a_e^\dagger(x) \partial_x a_e(x) + V(a_e^\dagger(0) \sum_{i=1}^N \sigma_{i-} + H.c.) + \sum_{i=1}^N (\Omega_i - i\gamma_i/2) a_{ei}^\dagger a_{ei}$ and $\mathcal{H}_o = -iv_g \int dx a_o^\dagger(x) \partial_x a_o(x)$. We set $v_g = 1$.

Two different 2LAs: We first calculate single- and two-photon transport in two different 2LAs, and next we generalize the results for many identical 2LAs. A single-photon scattering state $|k\rangle$ of a photon with energy $E_k = k$ is given by

$$|k\rangle = \int dx \{ A_1(g_k(x)a_e^\dagger(x) + \delta(x)e_{k,1}\sigma_{1+} + \delta(x)e_{k,2}\sigma_{2+}) + B_1 h_k(x)a_o^\dagger(x) \} |\emptyset\rangle, \quad (2)$$

where $g_k(x)$ ($h_k(x)$) is an amplitude of a single photon in the even (odd) mode while $e_{k,i}$ is an amplitude of the excited i th atom with $i = 1, 2$. Here A_1 and B_1 satisfy initial conditions of incident photon, i.e., photon is incoming in the left- or the right-moving channel. $|\emptyset\rangle$ represents vacuum state, i.e., atoms are in the ground state and zero photon in the waveguide. We derive the amplitudes in Eq.2 using the single-photon Schrödinger equation. The amplitudes are $e_{k,1} = V(E_k - \tilde{\Omega}_2)/(\sqrt{2\pi}\Xi)$, $e_{k,2} = V(E_k - \tilde{\Omega}_1)/(\sqrt{2\pi}\Xi)$, $g_k(x) = (\theta(-x) + t_2(k)\theta(x))e^{ikx}/\sqrt{2\pi}$ and $h_k(x) = e^{ikx}/\sqrt{2\pi}$ where $\Xi = (E_k - \tilde{\Omega}_1 + i\Gamma/2)(E_k - \tilde{\Omega}_2 + i\Gamma/2) + \Gamma^2/4$, $\Gamma = V^2$ and

$$t_2(k) = [(E_k - \tilde{\Omega}_1 - i\frac{\Gamma}{2})(E_k - \tilde{\Omega}_2 - i\frac{\Gamma}{2}) + \frac{\Gamma^2}{4}]/\Xi. \quad (3)$$

The single-photon transmission coefficient $T_k = |t_2(k) + 1|^2/4$, and reflection coefficient $R_k = |t_2(k) - 1|^2/4$. We plot T_k and R_k in Fig.3. When two atoms are identical, $\Omega_1 = \Omega_2 = \Omega$ and $\gamma_1 = \gamma_2 = \gamma$ we find excitation




FIG. 3. Single-photon transmission T_k (blue full line) and reflection R_k (red dash line) coefficients with incident photon energy E_k for two two-level atoms. The parameters are, (a) $\Omega_1 = \Omega_2 = 0.5$, $\gamma_1 = \gamma_2 = 0.01$, (b),(c),(d) $\Omega_1 = 0.45$, $\Omega_2 = 0.55$, and (b) $\gamma_1 = \gamma_2 = 0.01$, (c) $\gamma_1 = \gamma_2 = 0.0$, (d) $\gamma_1 = 0.01$, $\gamma_2 = 0.0$. In all plots, $\Gamma = 0.04$.

amplitude of any atom, $e_2(k) = \frac{1}{\sqrt{2\pi}} \frac{V}{E_k - \Omega + i\gamma/2 + i\Gamma}$ and $t_2(k) \equiv \tau_2(k) = \frac{E_k - \Omega + i\gamma/2 - i\Gamma}{E_k - \Omega + i\gamma/2 + i\Gamma}$. For two identical atoms $T_k = 0$ and $R_k = 1$ at single photon resonance, i.e., $E_k = \Omega$ and $\gamma = 0$. T_k is unity and R_k is zero for two different 2LAs at $E_k = (\Omega_1 + \Omega_2)/2$ and $\gamma_1 = \gamma_2 = 0$. R_k is unity and T_k is zero for two different 2LAs when $E_k = \Omega_1$, $\gamma_1 = 0$ and/or $E_k = \Omega_2$, $\gamma_2 = 0$.

The general two-photon scattering state of two incident photons with energy $E_{\mathbf{k}} = E_{k_1} + E_{k_2} = k_1 + k_2$ has the form

$$|k_1, k_2\rangle = \int dx_1 dx_2 \left[A_2 \left\{ g(x_1, x_2) \frac{1}{\sqrt{2}} a_e^\dagger(x_1) a_e^\dagger(x_2) + e_1(x_1) \delta(x_2) a_e^\dagger(x_1) \sigma_{1+} + e_2(x_1) \delta(x_2) a_e^\dagger(x_1) \sigma_{2+} + e_3 \delta(x_1) \delta(x_2) \sigma_{1+} \sigma_{2+} \right\} + B_2 \left\{ j(x_1; x_2) a_e^\dagger(x_1) a_o^\dagger(x_2) + f_1(x_1) \delta(x_2) a_o^\dagger(x_1) \sigma_{1+} + f_2(x_1) \delta(x_2) a_o^\dagger(x_1) \sigma_{2+} \right\} + C_2 h(x_1, x_2) \frac{1}{\sqrt{2}} a_o^\dagger(x_1) a_o^\dagger(x_2) \right] |\emptyset\rangle, \quad (4)$$

where the physical meaning of various amplitudes is obvious from the definition. The coefficients, A_2, B_2 and C_2 determine initial conditions of incident photons. We find these amplitudes from two-photon Schrödinger equation with a choice of incoming two-photon state using the method of Ref.[18]. Let us call $\tilde{\Omega}_1 - \tilde{\Omega}_2 = \Delta$, $\sqrt{\Delta^2 - \Gamma^2} = \beta$ and $\lambda_{\pm} = E_{\mathbf{k}} - (\tilde{\Omega}_1 + \tilde{\Omega}_2)/2 + i\Gamma/2 \pm \beta/2$. We find

$$e_1(x) = \left(\frac{i(\Delta + \beta)}{\Gamma} c_1 e^{i\lambda_- x} + \frac{i(\Delta - \beta)}{\Gamma} c_2 e^{i\lambda_+ x} \right) \theta(x) + (e_{k_1,1} g_{k_2}(x) + e_{k_2,1} g_{k_1}(x)), \quad (5)$$

$$e_2(x) = (c_1 e^{i\lambda_- x} + c_2 e^{i\lambda_+ x}) \theta(x) + (e_{k_1,2} g_{k_2}(x) + e_{k_2,2} g_{k_1}(x)), \quad (6)$$

and $f_i(x) = e_{k_1,i} h_{k_2}(x) + e_{k_2,i} h_{k_1}(x)$, with $i = 1, 2$, and

$e_3 = i(t_2(k_1) + t_2(k_2) - 2)/(2\pi(E_{\mathbf{k}} - \tilde{\Omega}_1 - \tilde{\Omega}_2 + i\Gamma))$. Here

$$\begin{aligned} c_1 &= \frac{-i\Gamma - \Delta + \beta}{4\pi\beta} i(\Gamma\varepsilon_{k_1}\eta_{k_2} + \Gamma\varepsilon_{k_2}\eta_{k_1} - 2\pi V e_3), \\ c_2 &= \frac{i\Gamma + \Delta + \beta}{4\pi\beta} i(\Gamma\varsigma_{k_1}\eta_{k_2} + \Gamma\varsigma_{k_2}\eta_{k_1} - 2\pi V e_3), \end{aligned} \quad (7)$$

and $\eta_k = i(t_2(k) - 1)/\Gamma$, $\varepsilon_k = V/(k - (\tilde{\Omega}_1 + \tilde{\Omega}_2)/2 + i\Gamma/2 - \beta/2)$, $\varsigma_k = V/(k - (\tilde{\Omega}_1 + \tilde{\Omega}_2)/2 + i\Gamma/2 + \beta/2)$. The amplitudes of two-photon wavefunction are

$$\begin{aligned} g(x_1, x_2) &= \frac{1}{\sqrt{2}}(g_{k_1}(x_1)g_{k_2}(x_2) + g_{k_2}(x_1)g_{k_1}(x_2)) \\ &+ \left[\theta(x_1 - x_2)\theta(x_2) \left(\frac{\Delta + \beta - i\Gamma}{\sqrt{2}V} c_1 e^{i\lambda_- x_1} e^{i(E_{\mathbf{k}} - \lambda_-)x_2} \right. \right. \\ &\left. \left. + \frac{\Delta - \beta - i\Gamma}{\sqrt{2}V} c_2 e^{i\lambda_+ x_1} e^{i(E_{\mathbf{k}} - \lambda_+)x_2} \right) + (x_1 \leftrightarrow x_2) \right], \end{aligned} \quad (8)$$

$$j(x_1; x_2) = (g_{k_1}(x_1)h_{k_2}(x_2) + g_{k_2}(x_1)h_{k_1}(x_2)), \quad (9)$$

$$h(x_1, x_2) = \frac{1}{\sqrt{2}}(h_{k_1}(x_1)h_{k_2}(x_2) + h_{k_2}(x_1)h_{k_1}(x_2)) \quad (10)$$

The part of the wavefunctions in Eqs.5,6,8 involving c_1, c_2 is generated because of inelastic photon scattering by the atoms, and is a signature of background fluorescence. This part is also responsible for photon-photon interactions created by resonant interactions of the light beam with the atoms. By integrating out the field operators of the 2LSs from Eq.4 we write down an asymptotic outgoing scattering state using the original right and left moving free photons as a combination of two transmitted $t_2(x_1, x_2)$, two reflected $r_2(x_1, x_2)$ and one transmitted plus one reflected photon $rt(x_1, x_2)$.

$$\begin{aligned} &\int dx_1 dx_2 \left[t_2(x_1, x_2) \frac{1}{\sqrt{2}} a_R^\dagger(x_1) a_R^\dagger(x_2) + r_2(x_1, x_2) \right. \\ &\left. \frac{1}{\sqrt{2}} a_L^\dagger(x_1) a_L^\dagger(x_2) + rt(x_1, x_2) a_R^\dagger(x_1) a_L^\dagger(x_2) \right] |\emptyset\rangle. \end{aligned} \quad (11)$$

Later we discuss nature of various parts of the two-photon wavefunction in Eq.11 of the two different 2LAs in comparison with that of a V-type 3LA.

Multiple identical 2LAs: For two identical 2LAs, the expression in Eq.8 of the two-photon wavefunction in the even modes reduces to

$$\begin{aligned} g_2(x_1, x_2) &= \frac{1}{\sqrt{2}}(g_{2,k_1}(x_1)g_{2,k_2}(x_2) + g_{2,k_2}(x_1)g_{2,k_1}(x_2)) \\ &- \frac{iV}{\sqrt{2}} \mathcal{C}_2 \left[e^{iE_{\mathbf{k}}x_2 - i(\Omega - i\gamma/2 - i\Gamma)(x_2 - x_1)} \theta(x_2 - x_1) \theta(x_1) \right. \\ &\left. (x_1 \leftrightarrow x_2) \right], \text{ where } g_{2,k}(x) = \frac{e^{ikx}}{\sqrt{2\pi}} (\theta(-x) + \tau_2(k)\theta(x)), \\ \mathcal{C}_2 &= 8iV e_2(k_1)e_2(k_2) - \frac{2\sqrt{2}i\Gamma(e_2(k_1) + e_2(k_2))}{\sqrt{\pi}(E_{\mathbf{k}} - 2\Omega + i\gamma + i\Gamma)}. \end{aligned} \quad (12)$$

Here $\mathcal{C}_2$ is a measure of the strength of photon-photon interactions. The form of the two-photon wavefunction in the even modes for multiple identical 2LAs is almost similar to the one in Eq.12 except the spread of two-photon bound state (part of the wavefunction with $\mathcal{C}_2$)




FIG. 4. Correlation of two transmitted photons $|t_2(x_1, x_2)|^2$ for multiple identical 2LSs with distance separation $x_1 - x_2$ between photons. The left figure is for two incident photons at single-photon resonance, $E_{k_1} = E_{k_2} = \Omega$. $|t_2(x_1, x_2)|^2$ is zero for even number of 2LSs for $E_{k_1} = E_{k_2} = \Omega$. The right figure is at a detuned incident energy, $E_{k_1} = E_{k_2} = 0.5\Omega$ and $\Omega = 0.45$. The number of TLS in right figure is 1 to 4 for top to bottom curves. In all plots, $\Gamma = 0.04$ and $\gamma = 0$.

over $x_1 - x_2$ reduces with increasing number of atoms. For example, the two-photon wavefunction in the even modes for three identical 2LAs is

$$\begin{aligned} g_3(x_1, x_2) &= \frac{1}{\sqrt{2}}(g_{3,k_1}(x_1)g_{3,k_2}(x_2) + g_{3,k_2}(x_1)g_{3,k_1}(x_2)) \\ &- \frac{iV}{\sqrt{2}} \mathcal{C}_3 \left[e^{iE_{\mathbf{k}}x_2 - i(\Omega - i\gamma/2 - 1.5i\Gamma)(x_2 - x_1)} \theta(x_2 - x_1) \theta(x_1) \right. \\ &\left. (x_1 \leftrightarrow x_2) \right], \text{ where } g_{3,k}(x) = \frac{e^{ikx}}{\sqrt{2\pi}} (\theta(-x) + \tau_3(k)\theta(x)), \\ \mathcal{C}_3 &= 18iV e_3(k_1)e_3(k_2) - \frac{6\sqrt{2}i\Gamma}{\sqrt{\pi}} \frac{e_3(k_1) + e_3(k_2)}{E_{\mathbf{k}} - 2\Omega + i\gamma + 2i\Gamma} \end{aligned} \quad (13)$$

and $e_3(k) = V/(\sqrt{2\pi}(E_k - \tilde{\Omega} + 1.5i\Gamma))$, $\tau_3(k) = (E_k - \tilde{\Omega} - 1.5i\Gamma)/(E_k - \tilde{\Omega} + 1.5i\Gamma)$. We find for four identical 2LAs,

$$\mathcal{C}_4 = 32iV e_4(k_1)e_4(k_2) - \frac{12\sqrt{2}i\Gamma}{\sqrt{\pi}} \frac{e_4(k_1) + e_4(k_2)}{E_{\mathbf{k}} - 2\tilde{\Omega} + 3i\Gamma} \quad (14)$$

where $e_4(k) = V/(\sqrt{2\pi}(E_k - \tilde{\Omega} + 2i\Gamma))$ and $\tau_4(k) = (E_k - \tilde{\Omega} - 2i\Gamma)/(E_k - \tilde{\Omega} + 2i\Gamma)$. At single photon resonance, i.e., $E_k = \Omega$ and $\gamma = 0$, we find $\mathcal{C}_i = 0$ for an even number of identical 2LAs, and the magnitude of $\mathcal{C}_i$ falls with an increasing odd number of identical 2LAs (see left plot in Fig.4). Two degenerate incident photons at single-photon resonance are reflected from an even number of identical 2LAs by a purely elastic scattering process, thus no inelastic two-photon bound-state is formed in this case. Of course, $\mathcal{C}_i$ is not exactly zero even for an even number of atoms in a more physical situations with a finite value of loss (i.e., $\gamma \neq 0$) and a slightly detuned incident energy from the single photon resonance. We plot correlations of two transmitted photons in multiple identical 2LAs for incident photons at single-photon resonance (left plot of Fig.4) or detuned from single-photon resonance (right plot of Fig.4). Two photon correlations fall monotonically with an increasing number of 2LAs for two detuned incident photons.

V-type three level atom (3LA): Next we consider a V -type 3LA as shown in Fig.2. The Hamiltonian of a single 3LA in a 1D waveguide with right and left-moving photon modes is exactly identical to Eq.1 with $N = 2$. The single-photon transport in a V -type 3LA is the same as in the two 2LAs with similar transition energies. This is because single photon can excite only one atomic transition of two 2LAs or a 3LA at a time. Two photons can simultaneously excite two atomic transitions of two independent 2LAs, but not two transitions of a V -type 3LA where only one transition can be excited at a time. This creates a difference in two-photon transport in two 2LAs and in a V -type 3LA with similar energy configuration.

The two-photon transport in a V -type 3LA in a 1D waveguide can be found easily from the previous calculation for two different 2LAs by setting $e_3 = 0$ in Eq.4, because two excited levels of a 3LA can not be excited simultaneously. We compare nature of two transmitted and two reflected photons from two 2LAs and a 3LA in a 1D waveguide for different parameters. For simplicity we consider $\gamma_1 = \gamma_2 = 0$ in the following comparison, though a loss ($\gamma_i \neq 0$) develops little different features. (i) When two transitions of two 2LAs or a 3LA are identical, $\Omega_1 = \Omega_2 = \Omega$, two degenerate incident photons at $E_{k_1} = E_{k_2} = \Omega$ are perfectly reflected from two identical 2LAs, and the correlations of two reflected photons $|r_2(x_1, x_2)|^2$ is constant (which is equal to $1/2\pi^2$) for any distance separation $x = x_1 - x_2$ between them (see Fig.5(a)). The corresponding wavefunction of two transmitted photons $|t_2(x_1, x_2)|^2$ is zero at all x . In contrast, two reflected photons at these parameters for a V -type 3LA is anti-bunched, (check Fig.5(c)) while two transmitted photons are bunched together, Fig.5(b). The anti-bunching of two reflected photons in a side-coupled V -type 3LA occurs because two photons can not be emitted simultaneously by the 3LA, and the reflected photons are solely due to emission from the 3LA. (ii) Two degenerate incident photons with energy equal to $(\Omega_1 + \Omega_2)/2$ are fully transmitted through two different 2LAs, and the correlations of two transmitted photons is constant (equal to $1/2\pi^2$) at any x . The corresponding wavefunction of two reflected photons is zero at all x . The nature of two transmitted and two reflected photons for a 3LA at these parameter sets is identical to the two different 2LAs. (iii) The correlation of two transmitted photons for two different 2LAs in a 1D waveguide is zero at all x when the incident photon energy $E_{k_1} = \Omega_1$ and $E_{k_2} = \Omega_2$. The correlation of two reflected photons for two different 2LAs at these parameters shows a cosine oscillation with x (see Fig.5(d)). The nature of two transmitted and two reflected photons for a 3LA at these parameter sets is different from the two different 2LAs. Two transmitted photons from a 3LA are bunched (see Fig.5(e)) while two reflected photons are anti-bunched. The correlation of two reflected photons from a 3LA at these parameters also oscillates with x (see Fig.5(f)). In all three above sets, the total energy of incident photons is $(\Omega_1 + \Omega_2)$, when the fluorescence




FIG. 5. Correlations of two transmitted $|t_2(x_1, x_2)|^2$ (middle column) and two reflected $|r_2(x_1, x_2)|^2$ (first and last columns) photons with distance separation $x_1 - x_2$ between photons. The first column is for two 2LAs, and the middle and last columns are for a V -type 3LA. The parameters are, $\Omega_1 = \Omega_2 = E_{k_1} = E_{k_2} = 0.5$ (first row) and $\Omega_1 = E_{k_1} = 0.45$, $\Omega_2 = E_{k_2} = 0.55$ (second row). In all plots, $\Gamma = 0.04$, $\gamma_1 = \gamma_2 = 0$.

vanishes completely in two 2LAs, it has been discussed recently [14, 19]. However, a loss $\gamma_i \neq 0$ from the 2LAs can create non-zero fluorescence. The nature of two transmitted or two reflected photons in a system of two 2LAs and in a V -type 3LA are identical when the single photon transmission amplitude (which is same in both the systems) is exactly one. Then the amplitude e_3 of simultaneous excitation of both the 2LAs is zero, and the system of two 2LAs behaves like a V -type 3LA. Away from this special parameter set (which can be achieved easily with a slightly detuned incident photons or incorporating a loss $\gamma_i \neq 0$) the nature of two transmitted or two reflected photons in a system of two 2LAs and in a V -type 3LA are very different. Therefore, two-photon scattering by a tightly focused weak light beam can be used to differentiate between two 2LAs and a V -type 3LA with similar transition energies.

The single photon transmission in an ensemble of four 2LAs is similar to that in an ensemble of two V -type 3LAs with similar level structures, for example, a 2LA ensemble where two 2LAs have transition energy Ω_1 and other two have Ω_2 , while each 3LA in a 3LA ensemble has two transition energies Ω_1 and Ω_2 . Now, two V -type 3LAs are saturated by two photons because two photons can simultaneously excite two transitions of two different 3LAs. Still, two-photon transport in four 2LAs is different from that in two 3LAs. This is because two photons can excite any two atoms of four 2LAs, thus there are total six amplitudes of exciting any two 2LAs, while two transitions from two different 3LAs can be excited by two photons, therefore there are total four options for two simultaneous atomic transitions in the 3LAs. It creates a difference between two-photon transport in four 2LAs and two-photon transport in two V -type 3LAs. Let us consider a system of four identical 2LAs for which

we have already calculated the inelastic scattering contribution (see Eq.14). The inelastic contribution in two-photon scattering from two identical V -type 3LAs with two same transition energies $\Omega_1 = \Omega_2 = \Omega$ is given by

$$\mathcal{C}_{2V} = 32iV e_4(k_1)e_4(k_2) - \frac{8\sqrt{2}i\Gamma}{\sqrt{\pi}} \frac{e_4(k_1) + e_4(k_2)}{E_{\mathbf{k}} - 2\tilde{\Omega} + 2i\Gamma} \quad (15)$$

where $e_4(k)$ and $\tau_4(k)$ are as before. Away from single-photon resonance, $\mathcal{C}_{2V} \neq \mathcal{C}_4$. The difference in two-photon scattering also persists between six 2LAs and

three V -type 3LAs. However, with an increasing number of atoms this difference becomes weaker. Single photon transport in a driven Λ type 3LA [12, 16] where one transition is coupled to photon modes and another transition is driven by a classical laser beam, is also similar to the two 2LAs in a 1D waveguide. We find that two-photon transport in a driven Λ type 3LA matches with that in a V -type 3LA in a 1D waveguide.

The support of the U.S. Department of Energy through LANL/LDRD Program for this work is gratefully acknowledged.

-
- [1] I. Gerhardt, G. Wrigge, P. Bushev, G. Zumofen, M. Agio, R. Pfab, and V. Sandoghdar, Phys. Rev. Lett. **98**, 033601 (2007).
 - [2] G. Zumofen, N. M. Mojarad, V. Sandoghdar and M. Agio, Phys. Rev. Lett. **101**, 180404 (2008).
 - [3] J. Hwang, M. Pototschnig, R. Lettow, G. Zumofen, A. Renn, S. Götzinger and V. Sandoghdar, Nature **460**, 76 (2009).
 - [4] G. Hetet, L. Slodicka, M. Hennrich, and R. Blatt, Phys. Rev. Lett. **107**, 133002 (2011).
 - [5] J.-T. Shen and S. Fan, Phys. Rev. Lett. **98**, 153003 (2007); Phys. Rev. A **76**, 062709 (2007).
 - [6] D. E. Chang, A. S. Sorensen, E. A. Demler, and M. D. Lukin, Nature Phys. **3**, 807 (2007).
 - [7] A. V. Akimov et. al., Nature **450**, 402 (2007).
 - [8] T. Shi and C. P. Sun, Phys. Rev. B **79**, 205111 (2009).
 - [9] D. Roy, Phys. Rev. B **81**, 155117 (2010); Phys. Rev. A **83**, 043823 (2011).
 - [10] J. Q. Liao and C. K. Law, Phys. Rev. A **82**, 053836 (2010).
 - [11] P. Longo, P. Schmitteckert, and K. Busch, Phys. Rev. Lett. **104**, 023602 (2010).
 - [12] D. Roy, Phys. Rev. Lett. **106**, 053601 (2011).
 - [13] H. Zheng, D. J. Gauthier, and H. U. Baranger, Phys. Rev. Lett. **107**, 223601 (2011).
 - [14] E. Rephaeli, S. E. Kocabas and S. Fan, Phys. Rev. A **84**, 063832 (2011).
 - [15] H. Zheng and H. U. Baranger, arXiv: 1206.4442 (2012).
 - [16] D. Witthaut and A. S. Sorensen, New J. Phys. **12**, 043052 (2010).
 - [17] H. J. Carmichael, *An Open Systems Approach to Quantum Optics* (Lecture Notes in Physics) (Berlin: Springer).
 - [18] D. Roy, arXiv : 1209.0399 (2012).
 - [19] A. Muthukrishnan, G. S. Agarwal, and M. O. Scully, Phys. Rev. Lett. **93**, 093002 (2004).